APTITUDE HEALTH®

EPICS

SCCHN in 2022 and Beyond

October 26, 2022

Copyright © 2022 Aptitude Health. All Rights Reserved. APTITUDE HEALTH® is a federally registered service mark of Aptitude Health Holdings, LLC

Report Contents

EPICS

Content	Slide
Meeting Snapshot	3
Faculty Panel	4
Meeting Agenda	5
Key Insights and Strategic Recommendations	6
Current Immune-Based Approaches	9
Targeted Therapies, Alone or in Combination	18
Immune Checkpoint Therapy Optimization	28
Immune Checkpoint Inhibitors: Relapsed/Metastatic	38
Other Approaches (including investigational IOs)	53

Meeting Snapshot

EPICS

EPICS

VIRTUAL CLOSED-DOOR ROUNDTABLE

DATE:
October 26, 2022



DISEASE-STATE AND DATA PRESENTATIONS
by key experts

INSIGHTS REPORT
including postmeeting analyses and actionable recommendations

PANEL: 6 key experts in SCCHN in 4 states in the US and 2 countries in Europe

SCCHN-SPECIFIC DISCUSSIONS
on the latest research updates, therapeutic advances, and their application in clinical decision-making

Panel Consisting of 4 US and 2 European SCCHN Experts

EPICS



CHAIR
Ezra Cohen, MD, FRCPSC, FASCO
University of California San Diego Health

Robert L. Ferris, MD, PhD
UPMC Hillman Cancer Center

Tanguy Seiwert, MD
Johns Hopkins School of Medicine

Maura Gillison, MD, PhD
MD Anderson Cancer Center

Kevin Harrington, MBBS, MRCP, FRCR, FRCP, PhD, DIC
Institute of Cancer Research

Andreas Dietz, MD, PhD
University Hospital of Leipzig

APTITUDE HEALTH®

Meeting Agenda

EPICS

Time	Topic	Speaker
11.00 AM – 11.05 AM	Welcome and Introductions	Ezra Cohen, MD, FRCPSC, FASCO
11.05 AM – 11.15 AM	Current Immune-Based Approaches • Neoadjuvant • First line	Robert Ferris, MD, PhD
11.15 AM – 11.35 AM	Discussion	
11.35 AM – 11.40 AM	Key Takeaways	
11.40 AM – 11.50 AM	Targeted Therapies, Alone or in Combination	Maura Gillison, MD, PhD
11.50 AM – 12.10 PM	Discussion	
12.10 PM – 12.15 PM	Key Takeaways	
12.15 PM – 12.25 PM	Immune Checkpoint Therapy Optimization	Tanguy Seiwert, MD
12.25 PM – 12.40 PM	Discussion	
12.40 PM – 12.45 PM	Key Takeaways	
12.45 PM – 12.55 PM	Immune Checkpoint Inhibitors: Relapsed/Metastatic	Kevin Harrington, MBBS, MRCP, FRCR, FRCP, PhD, DIC
12.55 PM – 1.20 PM	Discussion	
1.20 PM – 1.25 PM	Key Takeaways	
1.25 PM – 1.35 PM	Other Approaches (including investigational IOs)	Andreas Dietz, MD, PhD
1.35 PM – 1.50 PM	Discussion	
1.50 PM – 1.55 PM	Key Takeaways	
1.55 PM – 2.00 PM	Meeting Wrap-up	Ezra Cohen, MD, FRCPSC, FASCO

APTITUDE HEALTH®

EPICS

Current Immune-Based Approaches

Robert Ferris, MD, PhD

UPMC Hillman Cancer Center

Abstracts Covered: Current Immune-Based Approaches

EPICS

- Neoadjuvant
  - 653O – Neoadjuvant nivolumab (N) before chemoradiation (CRT) in high-risk HPV driven

Primary results of the phase III KEYNOTE-412 study: Pembrolizumab (pembro) with chemoradiation therapy (CRT) vs placebo plus CRT for locally advanced (LA) head and neck squamous cell carcinoma (HNSCC)

Machiels et al. ESMO 2022. LBA5

EPICS

STUDY POPULATION (N = 804)

> Phase III trial in treatment-naive, unresected, LA HNSCC

PRIMARY ENDPOINT

Event-Free Survival, ITT Population

Data cutoff date: May 31, 2022

Phase II study of cemiplimab in patients with advanced cutaneous squamous cell carcinoma (CSCC): Final analysis from EMPOWER-CSCC-1 groups 1, 2 and 3

Migden et al. ESMO 2022. 814P

EPICS

STUDY POPULATION (N = 193)

> Open-label, nonrandomized, multicenter, international phase II study in

ENDPOINTS

	Group 1	Group 2	Group 3

metastatic cutaneous squamous cell carcinoma; Q2W, every 2 weeks; Q3W, every 3 weeks.

Effects of durvalumab-metformin on head and neck squamous cell carcinoma (HNSCC)bas revealed by digital spatial profiling (DSP)

Riordan et al. ESMO 2022. 689P

EPICS

STUDY POPULATION (n = 8)

> Window of opportunity trial for durva + met in SCCHN

PRIMARY ENDPOINT

Nivolumab (Nivo) and ipilimumab (Ipi) combined with radiotherapy (RT) in patients (pts) with locally advanced squamous cell carcinoma of the head and neck (LA SCCHN): Updated results on efficacy and correlative analysis

EPICS

Vathiotis et al. ESMO 2022. 672P

STUDY POPULATION (N = 24 [of whom 14 HPV positive])

> Patients with newly diagnosed stage IVA–B SCCHN of the oropharynx,

EFFICACY ENDPOINTS

APTITUDE HEALTH®

EPICS

Current Immune-Based Approaches

Key Takeaways

Key Takeaways: Current Immune-Based Approaches

EPICS

- We continue to see data in SCCHN, but the standard of care has not changed

Discussion Highlight: Current Immune-Based Approaches

EPICS

RECENT UPDATES AND ADVANCES IN IMMUNOTHERAPY FOR SCCHN TREATMENT

Of the studies presented, KEYNOTE-412 most impressed the panel of experts, but it raised several questions

randomized phase 3 study – (ICML) Zhang 408

- Experts believe the resolution of molecular phenotype is possible in real time, and these vulnerabilities can potentially be targeted

Promising tolerability and efficacy results from dose-escalation in an ongoing phase I/II study of tinostamustine with pritelutamab, nivolumab for relapsed/refractory B-cell NHL – (EHA) Ghosh #6222

- The regimen is seen as effective, exciting, safe, and broadly applicable to many patients

Brexucabtagene (brexucel) monotherapy for elderly/unfit patients with first-line DLBCL continues to show promising safety and efficacy with durable complete responses – (EHA) Chavez-Ai #EP1501

- This approach is seen as a great solution for a patient population in whom giving immunochemotherapy is difficult. It is viewed as effective and safe

FIRST-MIND: a phase 3b, open-label, randomized study to assess safety of telisotuzumab or telisotuzumab + lenalidomide in addition to R-CHOP in patients with newly diagnosed DLBCL – (EHA) Beriada #EP1506

- Experts believe the combination of telisotuzumab + lenalidomide with R-CHOP is safe. However, they would like to see phase III data to confirm its activity in this setting

Long-term analyses from L-MIND, a phase II study of telisotuzumab plus lenalidomide (len) in patients with R/R DLBCL – (ICML) Duell #08

- The L-MIND regimen is seen as useful in the specific patient population with refractory disease. It was noted to be effective, very safe, and well tolerated. Some of the responses seen were clearly very durable

APTITUDE HEALTH®

EPICS

Targeted Therapies, Alone or in Combination

Maura Gillison, MD, PhD

MD Anderson Cancer Center

Abstracts Covered: Targeted Therapies, Alone or in Combination

EPICS

- LBA33 – 5-year overall survival (OS) in patients (pts) with locally advanced squamous cell carcinoma of

5-year overall survival (OS) in patients (pts) with locally advanced squamous cell carcinoma of the head and neck (LA SCCHN) treated with xevinapant + chemoradiotherapy (CRT) vs placebo + CRT in a randomized, phase II study (1/2)

Bourhis et al. ESMO 2022. LBA33

EPICS

STUDY POPULATION (N = 96)

PRIMARY ENDPOINT

5-year overall survival (OS) in patients (pts) with locally advanced squamous cell carcinoma of the head and neck (LA SCCHN) treated with xevinapant + chemoradiotherapy (CRT) vs placebo + CRT in a randomized, phase II study (2/2)

EPICS

Bourhis et al. ESMO 2022. LBA33

Postoperative chemoradiation in combination with cetuximab followed by 6-months cetuximab maintenance in squamous cell carcinoma of the head and neck with high risk of locoregional recurrence

EPICS

Tamaskovics et al. ESMO 2022. 671P

Neoadjuvant chemotarget therapy with cetuximab, docetaxel, and cisplatin in locally advanced resectable oral/oropharygeal SCC: A multicenter phase III trial (Eagle trial)

Zhong et al. ESMO 2022 669P

EPICS

Real-world evidence of first-line cetuximab (CX) plus paclitaxel (PX) in recurrent or metastatic squamous cell carcinoma of the head and neck (SCCHN)

Cirauqui Cirauqui et al. ESMO 2022. 678P

EPICS

APTITUDE HEALTH®

EPICS

Targeted Therapies, Alone or in Combination

Key Takeaways

Key Takeaways: Targeted Therapies, Alone or in Combination

EPICS

> Experts are excited by the results from the phase II clinical trial with xevinapant.

Discussion Highlight: Targeted Therapies, Alone or in Combination

EPICS

EXPERTS WERE MOST IMPRESSED BY THE PHASE II RESULTS WITH

APTITUDE HEALTH®

APTITUDE HEALTH®

EPICS

Immune Checkpoint Therapy Optimization

Tanguy Seiwert, MD

Johns Hopkins School of Medicine

Abstracts Covered: Immune Checkpoint Therapy Optimization

EPICS

> 676P – ATHENA: A multicenter phase II of atezolizumab (A) and bevacizumab (B)

ATHENA: A multicenter phase II of atezolizumab and bevacizumab in patients with recurrent or metastatic squamous-cell carcinoma of the head and neck – The HPV-negative cohort

Fayette et al. ESMO 2022 676P

EPICS

APTITUDE HEALTH®

Evaluation of the DNA damage response network as predictor of nivolumab efficacy in head and neck squamous cell carcinoma

Economopoulou et al. ESMO 2022 661MO

EPICS

STUDY POPULATION (N = 76)

DATA

APTITUDE HEALTH®

OPTIM: A randomized phase II study on the OPTimization of IMmunotherapy in SCC of the head and neck – AIO-KHT-0117

Gruenwald et al. ESMO 2022 681P

EPICS

STUDY POPULATION (N = 154)

PRIMARY ENDPOINT

Next-generation sequencing (NGS) helps predict response to immunotherapy in recurrent or metastatic head and neck SCC patients

Mirallas et al. ESMO 2022 686P

EPICS

STUDY POPULATION (N = 96)

STUDY DATA

APTITUDE HEALTH®

APTITUDE HEALTH®

EPICS

Immune Checkpoint Therapy Optimization

Key Takeaways

Key Takeaways: Immune Checkpoint Therapy Optimization

EPICS

> The combination of anti–PD(L)-1 and anti-VEGF continues to be explored

Discussion Highlight: Immune Checkpoint Therapy Optimization (1/2)

EPICS

THE NEED FOR BIOMARKERS IS CLEAR. EXPERTS EAGERLY AWAIT A WAY TO MAXIMIZE TREATMENT EFFICACY USING A BIOMARKER-GUIDED APPROACH

APTITUDE HEALTH®

Discussion Highlight: Immune Checkpoint Therapy Optimization (2/2)

EPICS

THE BEST SEQUENCING OF IMMUNOTHERAPY FOR THE TREATMENT OF SCCHN IS STILL BEING INVESTIGATED

Single-agent nivolumab randomized to ipilimumab-nivolumab vs docetaxel is an evidence-free

“

Dr Harrington:

APTITUDE HEALTH®

EPICS

Immune Checkpoint Inhibitors: Relapsed/Metastatic

Kevin Harrington, MBBS, MRCP, FRCR, FRCP, PhD, DIC

Institute of Cancer Research

Abstracts Covered: Immune Checkpoint Inhibitors in R/M SCCHN

EPICS

- 658MO – Pembrolizumab (pembro) vs standard-of-care (SOC) in previously treated recurrent/metastatic

Pembrolizumab (pembro) vs standard-of-care (SOC) in previously treated recurrent/metastatic (R/M) head and neck squamous cell carcinoma (HNSCC): 6-year follow-up of KEYNOTE-040 (1/4)

EPICS

Soulieres et al. ESMO 2022. 658MO

Pembrolizumab (pembro) vs standard-of-care (SOC) in previously treated recurrent/metastatic (R/M) head and neck squamous cell carcinoma (HNSCC): 6-year follow-up of KEYNOTE-040 (2/4)

Soulieres et al. ESMO 2022. 658MO



Data cutoff date: February 3, 2022.

Soulières D, et al. ESMO 2022. Oral presentation.

Pembrolizumab (pembro) vs standard-of-care (SOC) in previously treated recurrent/metastatic (R/M) head and neck squamous cell carcinoma (HNSCC): 6-year follow-up of KEYNOTE-040 (3/4)

Soulieres et al. ESMO 2022. 658MO

Data cutoff date: February 3, 2022.

BICR, blinded independent central review; CI, confidence interval; CR, complete response; ITT, intent-to-treat; NA, not available; NE, non-estimable; ORR, objective response rate; PD, partial disease; PR, partial response; RECIST, Response Evaluation Criteria in Solid Tumors; SD, stable disease; SOC, standard of care.

Soulières D, et al. ESMO 2022. Oral presentation.

Pembrolizumab (pembro) vs standard-of-care (SOC) in previously treated recurrent/metastatic (R/M) head and neck squamous cell carcinoma (HNSCC): 6-year follow-up of KEYNOTE-040 (4/4)

EPICS

Soulieres et al. ESMO 2022. 658MO

Pembrolizumab with or without chemotherapy for first-line treatment of recurrent/metastatic (R/M) head and neck squamous cell carcinoma (HNSCC): 5-year results from KEYNOTE-048 (1/2)

EPICS

Tahara et al. ESMO 2022. 659MO



Pembrolizumab with or without chemotherapy for first-line treatment of recurrent/metastatic (R/M) head and neck squamous cell carcinoma (HNSCC): 5-year results from KEYNOTE-048 (2/2)

EPICS

Tahara et al. ESMO 2022. 659MO

APTITUDE HEALTH®

Pembrolizumab (pembro) + carboplatin (carbo) + paclitaxel (pacli) as first-line (1L) therapy in recurrent/metastatic (R/M) head and neck squamous cell carcinoma (HNSCC): Phase VI KEYNOTE-B10 study

EPICS

Dzienis et al. ESMO 2022. 651O

Effectiveness and quality-of-life (QoL) data from real-world study (ProNiHN) in patients (pts) with recurrent and/or metastatic squamous cell carcinoma of head and neck (R/M SCCHN) treated with nivolumab (nivo) in France

EPICS

Le Tourneau et al. ESMO 2022. 695P

HANNA: Real-world data of patients with recurrent and/or metastatic squamous cell carcinoma of the head and neck (R/M SCCHN), including first-line population, treated with nivolumab in Germany

EPICS

Langer et al. ESMO 2022. 680P



APTITUDE HEALTH®

EPICS

Immune Checkpoint Inhibitors: Relapsed/Metastatic

Key Takeaways

Key Takeaways: Immune Checkpoint Inhibitors: Relapsed/Metastatic

EPICS

- Recently released data are encouraging with regard to improving patient survival in
  - It is a strong prognostic factor for OS and EFS
- The first small-volume (≤3 mL) pull of bone marrow aspirate should be used for MRD quantification
- Flow-based and NGS-based MRD quantification have high correlation for ≥10^{-4} leukemia burden
  - Providers may choose which methodology works best in their practice environment
  - Emerging evidence supports use of NGS with deeper sensitivity
- On the basis of current evidence, serial monitoring of MRD is recommended
- Additional studies are needed to contextualize MRD for different genotypes and to determine optimal, and cost-effective, testing frequency

Discussion Highlight: Immune Checkpoint Therapy Optimization (1/2)

EPICS

USE OF CHEMOTHERAPY IN COMBINATION WITH IMMUNOTHERAPY

- It is a strong prognostic factor for OS and EFS

- The first small-volume (≤3 mL) pull of bone marrow aspirate should be used for MRD quantification

- Flow-based and NGS-based MRD quantification have high correlation for ≥10^{-4} leukemia burden
  - Providers may choose which methodology works best in their practice environment
  - Emerging evidence supports use of NGS with deeper sensitivity

- On the basis of current evidence, serial monitoring of MRD is recommended

- Additional studies are needed to contextualize MRD for different genotypes and to determine optimal, and cost-effective, testing frequency

Discussion Highlight: Immune Checkpoint Therapy Optimization (2/2)

EPICS

PEMBROLIZUMAB

- It is a strong prognostic factor for OS and EFS

- The first small volume (≤3 mL) pull of bone marrow aspirate should be used for MRD quantification

- Flow-based and NGS-based MRD quantification have high correlation for ≥10^{-4} leukemia burden
  - Providers may choose which methodology works best in their practice environment
  - Emerging evidence supports use of NGS with deeper sensitivity

- On the basis of current evidence, serial monitoring of MRD is recommended

- Additional studies are needed to contextualize MRD for different genotypes and to determine optimal, and cost-effective, testing frequency

APTITUDE HEALTH®

EPICS

Other Approaches (including investigational IOs)

Andreas Dietz, MD, PhD

University Hospital of Leipzig

Abstracts Covered: Other Approaches (including investigational IOs)

EPICS

- 652O – Randomized phase III VANCE study: Gemcitabine and carboplatin (GC)

Randomized phase III VANCE study: Gemcitabine and carboplatin (GC) followed by Epstein Barr virus-specific autologous cytotoxic T lymphocytes (EBV-CTL) vs the same chemotherapy as first-line treatment for advanced nasopharyngeal carcinoma (NPC)

Toh et al. ESMO 2022. 652O

EPICS

Phase I study of M6620 (VX-970, berzosertib) in combination with cisplatin and XRT in patients with locally advanced head and neck squamous cell carcinoma

EPICS

Bhatia et al. ESMO 2022. 656MO

Early response to leukocyte interleukin Injection (LI) immunotherapy extends overall survival (OS) in locally advanced primary squamous cell carcinoma of the head & neck (SCCHN): The IT-MATTERS study

Lavin et al. ESMO 2022. 690P

EPICS

OPTIM: A randomized phase II study on the OPTimization of IMmunotherapy in squamous carcinoma of the head and neck – AIO-KHT-0117

Gruenwald et al. ESMO 2022. 681P

EPICS

A phase II trial of pembrolizumab and cabozantinib in pts with recurrent metastatic head and neck SCC

Saba et al. ASCO 2022 6008

EPICS

APTITUDE HEALTH®

EPICS

Other Approaches (including investigational IOs)

Key Takeaways

Key Takeaways: Other Approaches (including investigational IOs)

EPICS

> The experts positively view attempts to test new MOAs to move the field forward

Discussion Highlight: Immune Checkpoint Therapy Optimization

EPICS

KINASE INHIBITORS

STUDY POPULATION

- MEDALIST study: MDS patients with ≤10% ring sideroblasts or ≤5% with *SF3B1* mutation, <5% blasts in bone marrow, non-del(5q) MDS, refractory/intolerant to ESAs, or naive, average RBC transfusion burden ≥2 U/8 weeks, no prior treatment with disease-modifying agents (eg, IMiDs, HMAIs); randomized to luspatercept (n = 153) or placebo (n = 76). This subanalysis: 96/153 patients who did not achieve RBC transfusion independence (TI) for ≥8 weeks by week 23 but continued treatment through week 48

OUTCOMES

- 56% (n = 71/96) of patients achieved RBC TI for ≥8 weeks
  - 24 hours, 1 patients achieved RBC TI for ≥24 weeks
  - Median time to achieving RBC TI for ≥8 weeks was 3 months

EXPERT CONCLUSIONS

Continuing luspatercept treatment beyond week 23 provides clinical benefit in late responders and decreases the transfusion load in patients